

Keladi Shivappa Nayaka University of Agricultural and Horticultural Sciences, Shivamogga, Karnataka

Gramin Krishi Mausam Sewa Project







Weather Forecast from 07.01.2023 to 11.01.2023

No. GKMSB/002/2023 Date: 06.01.2023

Weather Forecast Issued by the India Meteorological Department for Dakshina Kannada District for the coming five days – until 0830 hrs of 07.01.2023 to 11.01.2023

Sl.	Weather Deverators	Day-1	Day-2	Day-3	Day-4	Day-5	Past Week	Forecast
No	Weather Parameters	(07.01.2023)	(08.01.2023)	(09.01.2023)	(10.01.2023)	(11.01.2023)	Mean	Mean
1	Rainfall (mm)	0	0	0	0	0	0	0
2	Max Temp (°C)	34	34	34	34	34	34.7	34↓
3	Min Temp (°C)	22	22	22	22	21	19.9	22↑
4	Cloud cover (Okta)	1	1	1	1	2	0	1↑
5	Max. Relative Humidity (%)	80	80	82	82	82	79.6	81↑
6	Min. Relative Humidity (%)	39	37	35	34	35	44.8	36↓
7	Wind Speed (Kmph)	1	1	1	1	1	5.2	1↓
8	Wind Direction (°)	68	68	68	68	71	234	69↓
9	Forecast Warning	DRY	DRY	DRY	DRY	DRY		
10	Chance of Occurrence (%)	0	0	0	0	0		

As per the extended range rainfall forecast given by IMD, New Delhi and MC, Bangalore for coastal region of Karnataka including Dakshina Kannada district the rainfall will be Normal from 12.01.2023 to 18.01.2023

Summary of Weather Forecast for next Five days


As per the Weather forecast issued by the India Meteorological Department, Pune and Bengaluru for Dakshina Kannada district, **DRY weather followed by next five days.** The maximum temperature would be around 34°C and minimum temperature would be around 22°C to during next five days. The Morning Relative humidity (RH) would vary from 80-82 per cent, Evening Relative humidity (RH) would vary from 34-39 per cent and wind speed will be in the range of 1 km/hr.

Mobile Applications developed by India Meteorological Department for the benefit of Farming Community Farmers are suggested to download the app using below links

	Mausam:	Meghdoot:	Damini:
Andriod:	https://play.google.com/store/app	https://play.google.com/store/apps/detail	https://play.google.com/store/apps/det
	s/details?id=com.imd.masuam	s?id=com.aas.meghdoot	ails?id=com.lightening.live.damini
Apple	https://apps.apple.com/us/app/id1	https://apps.apple.com/in/app/meghdoot/	https://apps.apple.com/app/id1502385
OS:	<u>522893967</u>	<u>id1474048155</u>	<u>645</u>

Decadal Rainfall (mm) Comparison:

Year	30 years Mean	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022
Rainfall	3790.4	3628.1	2984.2	3325.3	2816.3	2383.4	2827.1	3000	3381.2	4118.4	4110.8	4110.8	3531.2
Rainy days	110	128	105	127	112	107	119	121	112	123	126	152	118



Weather based Agromet Advisories for Dakshina Kannada District

Horticulture Crops:

Horuculture Crop							
Crop	Operations/ Agromet Advisories						
General Advise	Farmers are advised to install Meghdoot, Mausam, Umang & Damini apps from mausam.imd.gov.in and be updated with weather events.						
Arecanut	To manage weeds in orchard spray Glyphosate (Amoniam salt)41%SL @ of						
(Weed	6ml/ltr or Glyphosate 54% SC @ 4ml/litre or Glyphosate 71% SC @ 6 gram per						
management)	litre.						
Arecanut	Farmers can Spray Hexaconazole or Propiconazole or Difenconazole 1ml/ltr to						
(Leaf Spot)	the infected leaves.						
Coconut	To manage weeds in orchard spray Glyphosate (Amoniam salt)41%SL @ of						
(Weed	6ml/ltr or Glyphosate 54% SC @ 4ml/litre or Glyphosate 71% SC @ 6 gram						
management)	per litre.						
Banana (General Advice)	 5 months After planting spray Banana special @5gm/ltr of water at an interval of 30days or one month.(For one sprayer spray solution add 1 lime and shampoo) Bunch Treatment: To get good yield & quality bunches treat the bunch with 10 gram of Urea,10 gm of Potassium Sulphate & 0.5kg of cowdung should be mixed with 100ml of water & tie to end of bunch 						
Black Pepper	Farmers can Mulch the root zone of a crop with dried leaves or crop residues to						
(Water	save the moisture or to avoid the loss of water from root zone. Farmers should						
management)	irrigate the crop upto 20litre per day per vien.						
Jasmine	For the Management of leaf spot disease, spray with Hexaconazole @ 1ml/litre						
(Leaf Spot)	of water.						
Cashew	Clean the infected place. After cleaning fill the infected part with						
(stem borer)	Chloropyriphos granules mixed with sand(1:1 ratio)						
Cashew (Tea Mosquito Bug)	Farmers should spray Monocrotophos @ of 1.7ml/ltr or Lamdacyalothrin @ 0.5ml/ltr.						
Water Melon (General Advise)	Right time to take sowing of Watermelon. Farmers are advised to go for seed treatment with Bavistin before sowing.						

Cereals and Pulses :

Agronomic	After the harvest of paddy farmers can go for sowing of Dicot crops like Black
Advice	gram, Green gram, Beans and Green Manure seeds in the residual moisture.
Groundnut	• After the harvesting of Paddy, Groundnut can be selected as the second crop. Right
(Land	time to take Primary tillage.
preparation)	• After the primary tillage farmers can till the land with secondary tillage instrument
	Rotovator. After that based on the moisture availability farmers can take sowing.
Groundnut	Before the sowing of Groundnut seeds should be treated with Rhizobium (1kg of
(Seed treatment)	seeds should be treated with 10gm of Rhizobium & after that seeds should be
	shade dried for 02 hours).
Groundnut	To manage weeds soil should be sprayed with Alacholar 50EC @ 3ml/ltr
(Land	
preparation)	

Animal Husbandry:

Cattle	Animals should be tied at closed rooms or cattle sheds to protect them from cold.								
(General advise)	> Cattle-Animals should be vaccinated against FMD, Black Quarter								
	Enterotoxaemia if not vaccinated earlier.								
	➤ Balanced feed (Green fodder + dry fodder) should be provided to avoid Acidiosis.								

(This Agro Advisory Information is based on Weather forecast received from MC, Bengaluru & IMD, New Delhi)

Mr. Praveen K.M, Technical Officer, GKMS, ZAHRS, Brahmavar, Udupi. Dr. K.V. Sudhir Kamath, Nodal Officer, GKMS, ZAHRS, Brahmavar, Udupi.